P - 03

Serial No. 10/645,869 Amendment dated July 25, 2005 In Reply of Office Action dated March 24, 2005

Amendments to the Claims:

This listing of claims will replace all prior versions and listings of claims in the specification.

Listing of Claims:

1(currently amended). A method for treating a human male subject having erectile dysfunction and for diagnosing diagnosed with penile atherosclerosis as a primary cause for the erectile dysfunction, the method producing an erection suitable for sexual intercourse, comprising:

- a) applying momentary vacuum therapy to the penis of the human male subject for reducing penile atherosclerosis:
- b) administering an erectile dysfunction medication to the human male subject for inducing an erection prior to intercourse;
- <u>c)</u> the human male subject being a subject for whom a PDE-5 inhibitor alone has never induced an erection suitable for sexual intercourse;

- d) the erectile dysfunction medication being a PDE-5 inhibitor; and
- e) the method being performed without the use of a constricting ring on the penis of the human male subject.
 - 2(cancelled).
 - 3(cancelled).
 - 4 (cancelled).

5(currently amended). A method as in claim $\frac{1}{2}$, $\frac{2}{2}$, wherein:

- b)—-the orectile dysfunction-medication is a PDE-5 inhibitor.
 - 6(cancelled).
 - 7(cancelled).
 - 8 (cancelled).

BEST AVAILABLE COPY

9(currently amended). A method as in claim 1, 2, wherein:

- a) the human male subject is a subject for whom a combination pharmacotherapy medication alone has never induced an erection suitable for sexual intercourse; and
- b) an additional the erectile dysfunction medication is administered with the PDE-5 inhibitor to provide a combination pharmacotherapy medication.

10 (currently amended). A method as in claim $\frac{1}{2}$, wherein:

- a) the human male subject is a subject for whom a combination pharmacotherapy medication alone has become one of less effective and ineffective in inducing an erection suitable for sexual intercourse; and
- b) an additional the erectile dysfunction medication is administered with the PDE-5 inhibitor to provide a combination pharmacotherapy medication.
 - 11(currently amended). A method as in claim 1, 2, wherein:
 - a) the human male subject is a subject for whom a

combination pharmacotherapy medication alone has never induced an erection suitable for sexual intercourse without adverse side effects; and

b) an additional the erectile dysfunction medication is administered with the PDE-5 inhibitor to provide a combination pharmacotherapy medication.

12(currently amended). A method as in claim $\frac{1}{2}$, wherein:

a) — the ercetile dysfunction-medication-is a vasbdilator;

b) the human male subject is a subject for whom a vasodilator used alone is ineffective; and

a) --e) the momentary vacuum therapy and administration of the PDE-5 inhibitor vacodilator are used by the human male subject periodically and regularly for eventually achieving a natural erection without the use of the vasodilator and momentary vacuum therapy.

13(original). A method as in claim 1, wherein:

a) the momentary vacuum therapy is used by the human

male subject periodically and regularly for eventually achieving a natural erection without the use of the momentary vacuum therapy.

14 (cancelled).

15(currently amended). A method as in claim 1, 14, wherein:

a) the <u>PDE-5 inhibitor</u> erectile dysfunction medication is sildenafil citrate.

16 (cancelled).

17(new). A method for treating a human male subject having erectile dysfunction and diagnosed with penile atherosclerosis as a primary cause for the erectile dysfunction, the method producing an erection suitable for sexual intercourse, comprising:

- a) applying momentary vacuum therapy to the penis of the human male subject for reducing penile atherosclerosis;
 - b) administering an erectile dysfunction medication

to the human male subject for inducing an erection prior to intercourse;

- c) the human male subject being a subject for whom a vasodilator alone has never induced an erection suitable for sexual intercourse;
- d) the erectile dysfunction medication being a vasodilator; and
- e) the method being performed without the use of a constricting ring on the penis of the human male subject.
 - 18 (new). A method as in claim 17, wherein:
 - a) the vasodilator is alprostadil.